

Method for Analysis of Different Oligosacchiride Structures

Elena Kostadinova · Pavlina Dolashka ·
Stefka Kaloyanova · Ludmyla Velkova ·
Todor Deligeorgiev · Wolfgang Voelter · Ivan Petkov

Received: 24 January 2012 / Accepted: 27 June 2012
© Springer Science+Business Media, LLC 2012

Abstract In this study, an improved, rapid, high yield synthesis of *N,N'*-4,4'-bis(benzyl-2-boronic acid)-bipyridinium dibromide (*o*-BBV) is described. The obtained *o*-BBV is applied in a two-component saccharide sensing system (complex) where it serves as a fluorescence quencher and a saccharide receptor. This system was applied to different natural oligosaccharides isolated from molluscan *Rapana venosa* (RvH1-a) and arthropodan *Carcinus aestuarii* (CaeH) hemocyanins (Hcs) and cyclodextrins (CDs). The carbohydrate contents of both Hcs were calculated in our previous work to be 1,6 % and 7 % for CaeH and RvH1-a, respectively. We propose that the difference in fluorescence increase of the native CaeH and RvH1-a when titrating them with the complex is due to the fact that the carbohydrate content of CaeH is lower and the carbohydrate chains are buried in between the structural subunits of the native molecule, while the glycans of the functional unit RvH1-a are exposed on the surface of the molecule leading to a 4-fold fluorescence's intensity change.

Keywords Cyclodextrins · Hemocyanins · Fluorescence spectroscopy · *Rapana venosa*

Introduction

The immune system is one of the most commonly studied in which glycans and glycoproteins play an important physiological role. Therefore, saccharide sensing is a current topic of interest in molecular recognition chemistry in determining saccharide concentrations in physiological fluids and clarifying the functions of glycoproteins at biomembranes [1, 2]. There are several methods to identify oligosaccharide structures, as well as glycosylation patterns on cells, tissues and body fluids [3–5]. Although several types of saccharide receptors applying different recognition mechanisms are available, the most well-understood receptors are based on arylboronic acids [5–10] because of their favourable stability in water, and easy detectable UV-VIS and fluorescence signals. In contrast to boronic acid-based saccharide recognition systems consisting of a single detector, we provide in this communication a two-component sensing system [11] comprising an anionic fluorescent dye (1) and a boronic acid-appended cationic viologen (2) (Fig. 1) that serves as fluorescence quencher and saccharide receptor [12–15].

Cyclodextrins (CDs) are a family of cyclic oligosaccharides, composed of α -1,4-linked glucopyranose subunits [16–18]. CDs are produced from starch by enzymatic degradation. α -Cyclodextrin (α -CD), β -cyclodextrin (β -CD) and γ -cyclodextrin (γ -CDs) are the most common CD species, referred to as first generation parent CDs (six, seven and eight glucosyl units, respectively). Generally, CDs can form host-guest complexes with a large variety of solid, liquid, and gaseous organic compounds by a molecular inclusion phenomenon. This inner inclusion exerts a profound effect on the physicochemical properties of the guest


E. Kostadinova · S. Kaloyanova · T. Deligeorgiev · I. Petkov
The Sofia University, Faculty of chemistry,
1 James Bourchier Blvd.,
1164 Sofia, Bulgaria

P. Dolashka (✉) · L. Velkova
Institute of Organic Chemistry, Bulgarian Academy of Sciences,
G. Bonchev 9,
1113 Sofia, Bulgaria
e-mail: pda54@abv.bg

P. Dolashka
e-mail: dolashka@orgchm.bas.bg

W. Voelter
Interfaculty Institute of Biochemistry, University of Tübingen,
Hoppe-Seyler-Strasse 4,
72076 Tübingen, Germany

Fig. 1 Proposed mechanism of saccharide detection: saccharide-induced dissociation of ground-state complex results in fluorescence increase



molecules as they are temporarily locked or caged within the host cavity, giving rise to beneficial modifications on the guest molecule's properties (solubility, reactivity, volatility) [19]. Therefore, the native CD modifications are effective templates for generating wide ranges of molecular hosts [20] and are employed as carriers for biologically active substances [21], enzyme models [22], separating agents [23], catalysts [24], mass transfer promoters [25], additives in perfumes, cosmetics, aliments or food [26], environmental protection agents [27], or sensors for organic molecules [28].

Hemocyanins (Hcs) are high molecular mass oxygen-transporting proteins, freely dissolved in the hemolymph of several arthropods and molluscs. Arthropodan Hcs occur as hexamers or multiples of hexamers (up to 8×6-mers) of approximately 75 kDa subunits, each containing a Cu(I) pair able to reversibly bind dioxygen [29, 30]. Molluscan Hcs, in contrast, are decamers or di-decamers of approximately 350–450 kDa subunits, and the (di)decameric structure has the shape of a hollow cylinder [31, 32]. Each structural subunit is organized by seven or eight globular functional units (FUs) of approximately 50 kDa. These FUs contain one dioxygen-binding Cu(I) pair and are assigned by the letters a–g (h) starting from the N-terminus [31–34]. Beside the quaternary structure, there are also large differences in the carbohydrate content and monosaccharide composition of Hcs from arthropods and molluscs. The carbohydrate content of arthropodan Hcs is relatively low (0.1–2 %, w/w) [35], while the molluscan Hcs usually have a higher carbohydrate content (2–9 %, w/w) and may contain unusual monosaccharides [36–39].

Results and Discussion

New methods for analyzing glycans and providing information about the glycosylation patterns on cells, tissues and in body fluids, are needed for diagnostics and treatment, as

well as for facilitating studies on the effects of glycosylation. Therefore, we represent herein the application of a complex between an anionic fluorescent dye and a boronic acid-appended cationic viologen, serving as a fluorescence quencher and a saccharide receptor.

A literature survey revealed two main synthetic procedures to obtain *N,N'*-4,4'-bis(benzyl-2-boronic acid)-bipyridinium dibromide, which differ in used solvents and reaction temperatures. The described methods for quaternization of 4,4'-bipyridine with *o*-bromomethylphenylboronic acid were carried out in different solvents—in methanol at room temperature for 15 h [40] or in DMF by heating to 70 °C for 48 h [41]. The most frequently described and cited method applies stirring for 15 h at room temperature in an inert atmosphere, followed by a multistep and laborious isolation procedure [40]. In order to shorten and improve the synthetic procedure as well as to avoid the use of a solvent, we simply melted the starting materials and so the reaction temperature was determined by the melting points of the starting compounds. It was found that if the temperature of the reaction mixture is below 100 °C, this does not allow obtaining of a homogenous melt at low reaction rate. Heating of the reaction mixture to 100–110 °C for 20 min (Fig. 2) was found to be optimal, giving a final product in high purity and excellent yield by TLC monitoring. The structure was confirmed by elemental analysis and ¹H NMR spectroscopy.

The absorbance spectrum of pyranine shows only one maximum at 432 nm and two maxima were identified at 440 nm and 470 nm after titration with *o*-BBV (Fig. 3). For our fluorescence spectroscopic measurements, a solution of pyranine (1.33×10^{-6}) in 50 mM phosphate buffer, pH 7.5, was titrated with increasing amounts of the *o*-BBV and a decrease (4-fold) in fluorescence emission of pyranine ($\lambda_{em}=508$ nm, $\lambda_{ex}=460$ nm) was observed (Fig. 4). The fluorescence quantum yield (Φ) of pyranine in our instrumental setup in aqueous solution has a value of 0.89. After titration of pyranine with *o*-BBV, a complex formation has




Fig. 2 Synthesis of *o*-bromomethylphenylboronic acid (**2**) and *N,N'*-bis-(benzyl-2-boronic acid)-[4,4']bipyridinium dibromide (*o*-BBV, **3**)

occurred, which cause reduction of the pyranine's quantum yield up to 0.18.

In the proposed mechanism in Fig. 1, the electrostatic association of the pyranine and the quencher results in ground-state complex formation, facilitating electron transfer from the dye to the viologen, which leads to a decrease in fluorescence intensity [42, 43]. When different saccharides are added to the system, formation of two anionic boronate esters effectively neutralize the dicationic viologen, thus greatly diminishing its quenching efficiency, and an increase in the fluorescence intensity of the dye is observed. Fluorescence modulation is therefore directly correlated with saccharide concentration. Verification of this method includes titration of the above mentioned complex with three types of cyclodextrins, namely α -CD, β -CD and acetyl- β -CD, as well as two different types of hemocyanins from molluscs and arthropods. All of the above mentioned components contain carbohydrates with different structures.

The experiment was started with titration of the simplest samples, namely α -CD, β -CD and acetyl- β -CD with the above described complex and the effect was estimated by fluorescence spectroscopy. The interaction of the complex with the hydroxyl groups in α -CD is represented by the change of fluorescence intensity (arbitrary units, a.u.) as it is shown on Fig. 5. Increase in the concentration of the titration agent does not lead to a significant change in the fluorescence intensity (12 a.u.) of the pyranine and the quantum yield has slightly increased from 0.18 to 0.21. The same results were observed for the other two cyclodehtrins, β -CD and acetyl- β -CD (results are not shown). All three tested cyclodextrins have




Fig. 3 UV-absorption spectra of pyranine (1×10^{-5} M) (-----) and pyranine (1×10^{-5} M) with *o*-BBV (3×10^{-4} M) (—)

similar behaviour towards the used complex (Fig. 5, insert). This lack of significant changes in the fluorescence intensity could be partially explained with the structure of these cyclic carbohydrates (Fig. 6).

As a consequence of the 4C_1 conformation of the glucopyranose units, all secondary hydroxyl groups are situated on one of the two edges of the ring, whereas all the primary ones are placed on the other edge. The ring, in reality, is a conical cylinder, which is frequently characterized as a




Fig. 4 **a** Fluorescence emission spectra of pyranine (1.33×10^{-6} M in 50 mM phosphate buffer, pH 7.5, $\lambda_{\text{ex}}=460$ nm and $\lambda_{\text{em}}=508$ nm) with increasing concentrations (0.0 – 1.67×10^{-5} M) of *o*-BBV; **b** Relationship between the fluorescence intensity and the concentration of the titration agents




Fig. 5 **a** Fluorescence emission spectra of the complex (pyranine (1.33×10^{-6} M) in presence of *o*-BBV) upon addition of α -CD with increasing concentrations (0.0 – 8×10^{-5} M); **b** Relationship between the fluorescence intensity and the concentration of the titration agents: (— \blacklozenge) α -CD; (— \blacksquare) β -CD and (— \blacktriangle) acetyl- β -CD




Fig. 6 Common structures of α -CD with 6 Glu, β -CD with 7 Glu and γ -CD with 8 Glu

doughnut or wreath-shaped truncated cone. The cavity is lined by the hydrogen atoms and the glycosidic oxygen bridges, respectively. The C-2-OH group of one glucopyranoside unit can form a hydrogen bond with the C-3-OH group of the adjacent glucopyranose unit. In the CD molecule, a complete secondary belt is formed by these hydrogen bonds; therefore the β -CD forms a rather rigid structure. This intramolecular hydrogen bond formation is probably the explanation for the observation of very low fluorescence effect of all CDs. Moreover, the number of free hydroxyl groups is limited effectively neutralizing the dicationic viologen, resulting in very low fluorescence intensity of the pyranine.

Our new carbohydrate determination method was tested with glycoproteins, hemocyanins, with different amount and structure of the glycans. Molluscan and arthropodan hemocyanins are giant oxygen-transporting glycoproteins which have the same function, but quite different structure [32–34]. Arthropodan Hcs exhibit unusual assemblies of up to 48 structural subunits with molecular masses of 75 kDa. Hemocyanin from crab *Carcinus aestuarii* (CaeH) is constituted of three major and two minor structural subunits with molecular masses of 75 kDa [35, 43]. The glycosylation analyses of *Carcinus* hemocyanin were reported which showed that the carbohydrate of native CaeH represents only 1.6 % of protein mass [35]. Therefore, this hemocyanin was chosen to test the utility and reliability of our new carbohydrate determination method.

A change in the fluorescence intensity with only 30 a.u. and the quantum yield with 0.064 during titration of the complex with different concentrations of CaeH is shown on Fig. 7. These data are comparable with those of the cyclodextrins' fluorescence intensity changes and could be explained with the oligosaccharide's structure of CaeH:




Fig. 7 **a** Fluorescence emission spectra of the complex (pyranine (1.33×10^{-6} M) in presence of *o*-BBV) upon addition of increasing concentrations (0 – 0.47×10^{-4} M) of CaeH; **b** Relationship between the fluorescence intensity and the concentration of the titration agents: CaeH (—◆—) and RvH1-a (—■—)

Two short O-glycan sequences with suggested structures (N-Acetyl-O-NeuAc₂Gal₃GalNAc₂ and N-Acetyl-O-di-NeuAc₂Gal₂GalNAc₂) and one N-glycan sequence (SO₄Man₄GlcNAc₃) were identified by MALDI-MS analyses and enzymatic digestions [35]. Carbohydrate groups of these glycans most probably are buried in between the structural subunits of the global protein CaeH and interactions through van der Waals forces or hydrogen bonds may be involved, preventing the access of the complex to the hydroxyl groups.

In contrast to this result, titration of the functional unit RvH1-a of the molluscan hemocyanin *Rapana venosa* with the complex leads to 4-fold increase in arbitrary units (a.u.) of fluorescence intensity and 0.46 units in the quantum yield (Fig. 8). Hemocyanins of molluscs are high molecular mass glycoproteins (9,000 kDa) with a complex quaternary structure organized by ten structural subunits of RvH1 and RvH2, arranged by 8 FUs [31–33]. The oligosaccharide structures of both structural subunits and of some FUs are very well studied [38]. Therefore, the reactivity of the






Fig. 8 Fluorescence emission spectra of the complex of pyranine (1.33×10^{-6} M) in presence of *o*-BBV, upon addition of increasing concentrations (0.0 – 1.4×10^{-4} M) of RvH1-a

complex pyranine (1.33×10^{-6} in 50 mM phosphate buffer, pH 7.5) and *o*-BBV, was tested using RvH1-a, a functional unit of the structural subunit RvH1, the carbohydrate content of which is known [44]. Several techniques were applied, including capillary electrophoresis, matrix-assisted laser desorption ionization MS, and electrospray ionization MS in combination with glycosidase digestion, to determine a carbohydrate content for RvH1-a of 7 % and structures for its N-glycosidically-linked carbohydrates as follows :

Glycan 1:



Glycan 2:



Based on the models of several FUs from different molluscan Hcs, we suggest that both glycans are exposed on the surface of the molecule. Thus their hydroxyl groups are not involved in any interactions, such as van der Waals forces or hydrogen bonds and as a consequence they can be easily accessed by the complex. This leads to a 4-fold increase in fluorescence intensity.

Conclusion

According to these preliminary results we suggest that our invention provides a new tool to determine accessible free hydroxyl groups of carbohydrates attached to glycoproteins, glycolipids and proteoglycans which are not buried within the macromolecular structure or fixed by hydrogen bridges. This kind of information may be of enormous value studying carbohydrate-based interaction between biomolecules or cells. Studies in our laboratories are under way testing

oligosaccharides and glycoproteins of different structure to develop a more quantitative test system.

Materials and Methods

All starting materials and solvents were commercial products purchased from Sigma-Aldrich (Germany). 2-Bromomethylphenylboronic acid was synthesized by modification of the reported procedure [40]. Analytical samples of the reaction products were obtained by recrystallization in ethanol. All products were characterized and compared with reported data. ^1H NMR spectra were obtained on a Bruker Advance II 600 MHz instrument in CDCl_3 as solvent. Elemental analyses were performed on a Vario III apparatus. Melting points were determined on a Kofler apparatus and are uncorrected.

Synthesis of 2-Bromomethylphenylboronic Acid

Synthesis of 2-Bromomethylphenylboronic acid was carried out as shown in Fig. 2. To a solution of 2-methylphenylboronic acid (**1**) (1 g, 0.0073 mol) in CCl_4 (20 ml) was added dropwise a solution of Br_2 (1.2 g, 0.0075 mol) in CCl_4 (3 ml). The reaction mixture was stirred and irradiated with an UV lamp for 1 h at room temperature and then refluxed for 2 h. The progress of the reaction was monitored by TLC (Merck F 254 silica gel; dichloromethane). After cooling, the resulting precipitate was filtered off from the reaction mixture, the filtrate concentrated under reduced pressure to give a yellowish-brown oil (**2**), which became solid after few days of storage in a refrigerator. The product was used without further purification. Yield 82 %. m.p. 156–159 °C (lit. m.p. 158–162 °C) [40].

Synthesis of N,N'-Bis-(Benzyl-2-Boronic Acid)-[4,4']bipyridinium Dibromide (o-BBV)

2-Bromomethylphenylboronic acid (**2**) (1.57 g, 0.0073 mol) and 4,4'-bipyridine (0.38 g, 0.0024 mol) were mixed together and heated at 100–110 °C for 20 min. The progress of the reaction was monitored by TLC (Merck F 254 silica gel; dichloromethane: methanol: acetic acid, 86:13:1). After cooling of the resulting yellow-brown melt to room temperature, 10 ml ethanol was added to dissolve the solid mass and then the solution was diluted with 20–30 ml diethyl ether. The product precipitated as yellow-brown solid, which was filtered off, washed with diethyl ether, and recrystallized in ethanol. Yield 69 %, m.p. (ethanol) 280–283 °C (decomposition).

^1H NMR (CDCl_3): δ 6.07 (s, 4H, NCH_2), 7.36–7.53 (m, 6H, ArH), 7.78 (d, 2H, ArH), 7.83 (d, 2H, ArH), 8.78 (d, 2H, ArH), 9.38 (d, 2H, ArH), 9.48 (d, 2H, ArH). Elemental

analysis calcd for $\text{C}_{24}\text{H}_{24}\text{B}_2\text{Br}_2\text{N}_2\text{O}_4$: C, 49.20; H, 4.13; N, 4.78. Found: C, 48.89; H, 3.92; N, 4.56.

Isolation of FU of Hemocyanins

The glycosylated functional unit RvH1-a was isolated via an anion-exchange chromatography column on a FPLC system, as described by Dolashka et al [34]. Native Hc from the crab *C. aestuarii* was prepared from the hemolymph obtained from the dorsal lacuna of living animals collected in the lagoon of Venice as reported by Dolashka et al [35].

Fluorescence Measurements

Absorbance spectra were recorded in a Shimadzu spectrometer of (1×10^{-5} M) pyranine and after titration of pyranine with o-BBV (3×10^{-4} M). Fluorescence measurements were carried out on a spectrofluorimeter Jasco FP-6600. The emission spectra of pyranine (1.33×10^{-6}) in 50 mM phosphate buffer, pH 7.5, were measured between 460 and 650 nm, with excitation wavelength at 460 nm in a 1 cm quartz cuvette upon addition of samples with different concentrations. For Caeh and RvH1-a the concentration range was within 0 – 1×10^{-4} mol/l and for cyclodextrin 0 – 8×10^{-6} mol/l. All studies were carried out under ambient conditions (25 °C, in air).

Acknowledgments This work was supported by a research grant by the Bulgarian National Science Fund TK01-496/2009 and DFG-STE 1819/5-1/2012 (Germany). P. Dolashka and E. Kostadinova thank to German Academic Exchange Service (DAAD) for financing this study.

References

- Geddes CD, Lakowicz JR (2006) Topics in fluorescence spectroscopy. Springer, New York, pp 344–350
- Suzuki I, Yamauchi A, Sakashita Y, Hirose K, Miura T, Hayashita T (2007) Fluorescence response mechanism of D-glucose selectivity for supramolecular probes composed of phenylboronic-acid-modified β -cyclodextrin and styrylpyridinium dyes. Anal Sci 23:1167–1171
- James TD, Linnane P, Shinkai S (1996) Fluorescent saccharide receptors: a sweet solution to the design, assembly and evaluation of boronic acid derived PET sensors. Chem Commun 3:281–288
- Badugu R, Lakowicz JR, Geddes CD (2005) Boronic acid fluorescent sensors for monosaccharide signaling based on the 6-methoxyquinolinium heterocyclic nucleus: progress toward noninvasive and continuous glucose monitoring. Bioorg Med Chem 13:113–119
- Badugu R, Lakowicz JR, Geddes CD (2004) Noninvasive continuous monitoring of physiological glucose using a monosaccharide-sensing contact lens. Anal Chem 76:610–618
- Cao H, McGill T, Heagy MD (2004) Substituent effects on monoboronic acid sensors for saccharides based on N-phenyl-1,8-naphthalenedicarboximides. J Org Chem 69:2959–2966
- Cordes DB, Gamsey S, Sharrett Z, Miller A, Thoniyot P, Wessling RA, Singaram B (2005) The interaction of boronic acid-substituted viologens with pyranine: the effects of quencher charge on

- fluorescence quenching and glucose response. *Langmuir* 21:6540–6547
8. Zhang Y, Gao Z, Hardcastle K, Wang B, Zhang Y, Gao Z, Hardcastle K, Wang B (2006) Water-soluble fluorescent boronic acid compounds for saccharide sensing: substituent effects on their fluorescence properties. *Chem Eur J* 12:1377–1384
 9. Jiang S, Escobedo JO, Kim KK, Alptürk O, Samoei GF, Fakayode SO, Warner IM, Rusin O, Strongin RM (2006) Stereochemical and regiochemical trends in the selective detection of saccharides. *J Am Chem Soc* 128:12221–12228
 10. Tan W, Zhang D, Wg Z, Li C, Zhu D (2007) 4-(N, N dimethylamine)benzonitrile (DMABN) derivatives with boronic acid and boronate groups: new fluorescent sensors for saccharides and fluoride ion. *J Mater Chem* 17:1964–1968
 11. Camara JN, Suri JT, Cappuccio FE, Wessling RA, Singaram B (2002) Boronic acid substituted viologen based optical sugar sensors: modulated quenching with viologen as a method for monosaccharide detection. *Tetrahedron Lett* 43:1139–1141
 12. Arimori S, Ward CJ, James TD (2002) A d-glucose selective fluorescent assay. *Tetrahedron Lett* 43:303–305
 13. Wiskur SL, Ait-Haddou H, Lavigne JJ, Anslyn EV (2001) Teaching old indicators new tricks. *Acc Chem Res* 34:963–972
 14. Arimori S, Murakami H, Takeuchi M, Shinkai S (1995) Sugar-controlled association and photoinduced electron-transfer in boronic-acid-appended porphyrins. *Chem Commun* :961–962
 15. DiCesare N, Pinto MR, Schanze KS, Lakowicz JR (2002) Saccharide detection based on the amplified fluorescence quenching of a water-soluble poly(phenylene ethynylene) by a boronic acid functionalized benzyl viologen derivative. *Langmuir* 18:7785–7787
 16. Bender ML, Komiyama M (1978) *Cyclodextrin chemistry*. Springer, Berlin
 17. Szejtli J (1988) *Cyclodextrin technology*. Kluwer Academic Publishers, The Netherlands
 18. Szejtli J (1998) Introduction and general overview of cyclodextrin chemistry. *Chem Rev* 98:1743–1753
 19. Singh M, Sharma R, Banerjee UC (2002) Biotechnological applications of cyclodextrins. *Biotechnol Adv* 20:341–359
 20. Szejtli J (1982) *Cyclodextrins and their inclusion complexes*. Akademiai Kiado, Budapest
 21. Uekama K (2002) Recent aspects of pharmaceutical application of cyclodextrins. *J Incl Phenom Macrocycl Chem* 44:3–7
 22. Fernandez M, Frago A, Cao R, Villalonga R (2005) Stabilization of α -chymotrypsin by chemical modification with monoamine cyclodextrin. *Process Biochem* 40:2091–2094
 23. Lelievre F, Yan C, Zare RN, Gareil P (1996) Capillary electrochromatography: operating characteristics and enantiomeric separations. *J Chromatogr A* 723:145–156
 24. Bhosale SV (2007) b-Cyclodextrin as supramolecular catalyst in organic synthesis. *Synlett* 1:175–176
 25. Blach P, Landy D, Fourmentin S, Surpateanu G, Bricout H, Ponchel A, Hapiot F, Monflier E (2005) Sulfobutyl ether- β -cyclodextrins: promising supramolecular carriers for aqueous organometallic catalysis. *Adv Synth Catal* 347:1301–1307
 26. Szenté L, Szejtli J (2004) Cyclodextrins as food ingredients. *Trends Food Sci Technol* 15:137–142
 27. Fourmentin S, Outirite M, Blach P, Landy D, Ponchel A, Monflier E, Surpateanu G (2007) Solubilisation of chlorinated solvents by cyclodextrin derivatives: a study by static headspace gas chromatography and molecular modeling. *J Hazard Mater* 141:92–97
 28. Hattori K, Takeuchi T, Ogata M, Takanohashi A, Mikuni K, Nakanishi K, Imata H (2007) Detection of environmental chemicals by SPR assay using branched cyclodextrin as sensor ligand. *J Incl Phenom Macrocycl Chem* 57:339–342
 29. Burmester T (2001) Molecular evolution of the arthropod hemocyanin superfamily. *Mol Biol Evol* 18:184–195
 30. Dolashka-Angelova P, Dolashki A, Stevanovic S, Hristova R, Atanasov B, Nicolov P, Voelter W (2005) Structure and stability of arthropodan hemocyanin *Limulus polyphemus*. *Spectrochim Acta* 6:1207–1217
 31. Gatsogiannis Ch, Markl J (2009) Keyhole limpet hemocyanin: 9-Å CryoEM structure and molecular model of the KLH1 dodecamer reveal the interfaces and intricate topology of the 160 functional units. *J Mol Biol* 385:963–983
 32. Dolashka-Angelova P, Schwarz H, Dolashki A, Stevanovic S, Fecker M, Saeed M, Voelter W (2003) Oligomeric stability of *Rapana venosa* hemocyanin (RvH) and its structural subunits. *Biochim Biophys Acta* 1646:77–85
 33. Lieb B, Boissguérin V, Gebauer W, Markl J (2004) cDNA sequence, protein structure, and evolution of the single hemocyanin from *Aplysia californica*, an opisthobranch gastropod. *J Mol Evol* 59:536–545
 34. Dolashka-Angelova P, Stevanovic S, Dolashki A, Devreese B, Tzvetkova B, Voelter W, Van Beeumen J, Salvato B (2007) A challenging insight on the structural unit 1 of molluscan *Rapana venosa* hemocyanin. *Arch Biochem Biophys* 459:50–58
 35. Dolashka-Angelova P, Beltramini M, Dolashki A, Salvato B, Voelter W (2001) Carbohydrate composition of *carcinus aestuarii* hemocyanin. *Biophys* 389:153–158
 36. Velkova L, Dolashka P, Lieb B, Dolashki A, Voelter W, Van Beeumen J, Devreese B (2011) Glycan structures of the structural subunit (HtH1) of *Haliotis tuberculata* hemocyanin. *Glycoconj J* 28:385–395
 37. Wührer M, Robijn MLM, Koeleman CAM, Balog CIA, Geyer R, Deelder AM, Hokke CH (2004) A novel Gal(β 1-4)Gal(β 1-4)Fuc(α 1-6)-core modification attached to the proximal N-acetylglucosamine of keyhole limpet haemocyanin (KLH) N-glycans. *Biochem J* 378:625–632
 38. Dolashka P, Velkova L, Shishkov S, Kostova K, Dolashki A, Dimitrov I, Atanasov B, Devreese B, Voelter W, Beeumen V (2010) Glycan structures and antiviral effect of the structural subunit RvH2 of *Rapana* hemocyanin. *J Carbohydr Res* 345:2361–2367
 39. Lommerse JP, Thomas-Oates JE, Gielen C, Préaux G, Kamerling JP, Vliegthart JF (1997) Primary structure of 21 novel monomeric and dimeric N-linked carbohydrate chains from α -D-hemocyanin from *Helix pomatia*. *Eur J Biochem* 249:195–222
 40. Snyder HR, Reedy AJ, Lennarz WJ (1958) Synthesis of aromatic boronic acids. Aldehyde boronic acids and a boronic acid analog of tyrosine. *J Am Chem Soc* 80:835–838
 41. Gamsey S, Miller A, Olmstead M, Beavers C, Hirayama L, Pradhan S, Wessling R, Singaram B (2007) Boronic acid-based bipyridinium salts as a tunable receptors for monosaccharides and α -hydroxycarboxylates. *J Am Chem Soc* 129:1278–1286
 42. Cappuccio FE, Suri JT, Cordes DB, Wessling RA, Singaram B (2004) Evaluation of pyranine derivatives in boronic acid based saccharide sensing: significance of charge interaction between dye and quencher in solution and hydrogel. *J of Fluoresc* 14:521–533
 43. Dolashka-Angelova P, Dolashki A, Savvides SN, Hristova R, Van Beeumen J, Voelter W, Devreese B, Weser U, Di Muro P, Salvato B, Stevanovic S (2005) Structure of Hemocyanin subunit CaeSS2 of the Crustacean mediterranean crab *carcinus aestuarii*. *J Biochem* 138:303–312
 44. Dolashka-Angelova P, Beck A, Dolashki A, Beltramini M, Stevanovic S, Salvato B, Voelter W (2003) Characterization of the carbohydrate moieties of the functional unit RvH1-a of *Rapana venosa* haemocyanin using HPLC/electrospray ionization MS and glycosidase digestion. *Biochem J* 374:185–192